

FORM PTO-1390
(REV. 10-96)

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER
19424PC

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

08/913644

INTERNATIONAL APPLICATION NO.

PCT/US96/03649

INTERNATIONAL FILING DATE

18-Mar-1996

PRIORITY DATE CLAIMED

22-Mar-1995

TITLE OF INVENTION

DNA ENCODING HUMAN PAPILOMAVIRUS TYPE 18

APPLICANT(S) FOR DO/EO/US

KATHRYN J. HOFMANN, KATHRIN U. JANSEN, MICHAEL P. NEEPER, JOSEPH G. JOYCE, HUGH A. GEORGE

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(I).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

EXPRESS MAIL CERTIFICATE

DATE OF DEPOSIT September 17, 1997

EXPRESS MAIL NO. EM230313095US

I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS
BEING DEPOSITED WITH THE UNITED STATES POSTAL
SERVICE AS EXPRESS MAIL "POST OFFICE TO
ADDRESSEE" BEFORE 5 P.M. ON THE ABOVE DATE IN
AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER
FOR PATENTS, WASHINGTON, D.C. 20231.

MAILED BY [Signature]
DATE Sept. 17, 1997

Rec'd PCT/PTO 17 SEP 1997

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)	INTERNATIONAL APPLICATION NO. PCT/US96/03649	ATTORNEY'S DOCKET NUMBER 19424PC
---	---	-------------------------------------

17. <input type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Search Report has been prepared by the EPO or JPO.....\$910.00 International preliminary examination fee paid to USPTO (37 CFR 1.482).....\$700.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).....\$770.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$1,040.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4).....\$ 96.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	CALCULATIONS	PTO USE ONLY
	\$96.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	\$130.00	

Claims	Number Filed	Number Extra	Rate		
Total Claims	20	-20=	0	X \$22.00	\$0.00
Independent Claims	3	- 3=	0	X \$80.00	\$0.00
Multiple dependent claim(s) (if applicable)				+ \$260.00	\$0.00
TOTAL OF ABOVE CALCULATIONS				=	\$226.00
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).					
SUBTOTAL				=	\$226.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$0.00
TOTAL NATIONAL FEE				=	\$226.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +					
TOTAL FEES ENCLOSED				=	\$226.00
					Amount to be refunded
					charged

a. ☐ A check in the amount of \$0.00 to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 13-2755 in the amount of \$226.00 to cover the above fees. A duplicate copy of this sheet is enclosed.


c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any over-payment to the Deposit Account No. 13-2755. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Patent Department, RY60-30
 MERCK & CO., INC.
 P.O. Box 2000
 126 East Lincoln Avenue
 Rahway, New Jersey 07065

DATE: September 17, 1997


 SIGNATURE
 JOANNE M. GIESSER
 NAME
 32,838
 REGISTRATION NUMBER

10 Rec'd 7/27/97 17 SEP 1997

- 1 -

TITLE OF THE INVENTIONDNA ENCODING HUMAN PAPILLOMAVIRUS TYPE 18CROSS-RELATED TO OTHER APPLICATIONS

- 5 This is a continuation of U.S. Serial No. 08/408,669 filed March 22, 1995, now pending, and a continuation of U.S. Serial No. 08/409,122 filed March 22, 1995, now pending.

FIELD OF THE INVENTION

- 10 The present invention is directed to DNA molecules encoding purified human papillomavirus type 18 and derivatives thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

- 15 Figure 1 shows the HPV18 L1 nucleotide and deduced amino acid sequences.

Figure 2 is a list of amino acid variations within the L1 protein of HPV 18.

Figure 3 shows the HPV18 L2 nucleotide and deduced amino acid sequences.

- 20 Figure 4 shows an immunoblot of HPV18 L1 protein expressed in yeast.

Figure 5 shows an immunoblot of HPV18 L2 protein expressed in yeast.

- 25 Figure 6 is an electron micrograph of virus-like particles formed by HPV18 L1 protein expressed in yeast.

BACKGROUND OF THE INVENTION

- 30 Papillomavirus (PV) infections occur in a variety of animals, including humans, sheep, dogs, cats, rabbits, monkeys, snakes and cows. Papillomaviruses infect epithelial cells, generally inducing benign epithelial or fibroepithelial tumors at the site of infection. PV are species specific infective agents; a human papillomavirus does not infect a nonhuman animal.

- 2 -

Papillomaviruses may be classified into distinct groups based on the host that they infect. Human papillomaviruses (HPV) are further classified into more than 70 types based on DNA sequence homology. PV types appear to be type-specific immunogens in that a neutralizing immunity to infection by one type of papillomavirus does not confer immunity against another type of papillomavirus.

In humans, different HPV types cause distinct diseases. HPV types 1, 2, 3, 4, 7, 10 and 26-29 cause benign warts in both normal and immunocompromised individuals. HPV types 5, 8, 9, 12, 14, 15, 17, 19-25, 36 and 46-50 cause flat lesions in immunocompromised individuals. HPV types 6, 11, 34, 39, 41-44 and 51-55 cause benign condylomata of the genital or respiratory mucosa. HPV types 16 and 18 cause epithelial dysplasia of the genital mucosa and are associated with the majority of *in situ* and invasive carcinomas of the cervix, vagina, vulva and anal canal.

Papillomaviruses are small (50-60 nm), nonenveloped, icosahedral DNA viruses that encode for up to eight early and two late genes. The open reading frames (ORFs) of the virus genomes are designated E1 to E7 and L1 and L2, where "E" denotes early and "L" denotes late. L1 and L2 code for virus capsid proteins. The early (E) genes are associated with functions such as viral replication and cellular transformation.

The L1 protein is the major capsid protein and has a molecular weight of 55-60 kDa. The L2 protein is a minor capsid protein which has a predicted molecular weight of 55-60 kDa and an apparent molecular weight of 75-100 kDa as determined by polyacrylamide gel electrophoresis. Immunological data suggest that most of the L2 protein is internal to the L1 protein within the viral capsomere. The L1 ORF is highly conserved among different papillomaviruses. The L2 proteins are less conserved among different papillomaviruses.

The L1 and L2 genes have been identified as good targets for immunoprophylaxis. Studies in the cottontail rabbit papillomavirus (CRPV) and bovine papillomavirus (BPV) systems have shown that immunizations with the L1 and L2 proteins expressed in bacteria or by

- 3 -

using vaccinia vectors protected animals from viral infection. Expression of papillomavirus L1 genes in baculovirus expression systems or using vaccinia vectors resulted in the assembly of virus-like particles (VLP) which have been used to induce high-titered virus-neutralizing antibody responses that correlate with protection from viral challenge.

Following HPV type 16, HPV18 is the second most prevalent HPV type found in cervical carcinomas. HPV18 was detected in 5-20% of cervical cancer biopsies collected from various parts of the world (Ikenberg, H. 1990. Human papillomavirus DNA in invasive genital carcinomas. In Genital Papillomavirus Infections, G. Gross *et al.*, eds. p. 85-112). There appears to be a geographic dependence of infection with HPV18 since tumor biopsies from African and South American women harbor HPV18 more frequently than similar biopsies from European and North American women. The underlying reasons for these geographic differences are not known. The development of a vaccine against HPV18 infection becomes extremely relevant since HPV18 is also associated with more aggressively growing cancers and is rarely found in the milder precursor lesions, CIN I-II.

SUMMARY OF THE INVENTION

The present invention is directed to DNA molecules encoding purified human papillomavirus type 18 (HPV type 18; HPV18) and uses of the DNA molecules.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to DNA molecules encoding purified human papillomavirus type 18 (HPV type 18; HPV18) and derivatives thereof. Such derivatives include but are not limited to peptides and proteins encoded by the DNA, antibodies to the DNA or antibodies to the proteins encoded by the DNA, vaccines comprising the DNA or vaccines comprising proteins encoded by the DNA, immunological compositions comprising the DNA or the proteins encoded by the DNA, kits containing the DNA or RNA derived from the DNA or proteins encoded by the DNA.

- 4 -

Pharmaceutically useful compositions comprising the DNA or proteins encoded by the DNA may be formulated according to known methods such as by the admixture of a pharmaceutically acceptable carrier. Examples of such carriers and methods of formulation may be found in Remington's Pharmaceutical Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the DNA or protein or VLP. Such compositions may contain DNA or proteins or VLP derived from more than one type of HPV.

Therapeutic or diagnostic compositions of the invention are administered to an individual in amounts sufficient to treat or diagnose PV infections. The effective amount may vary according to a variety of factors such as the individual's condition, weight, sex and age. Other factors include the mode of administration. Generally, the compositions will be administered in dosages ranging from about 1 μ g to about 1 mg.

The pharmaceutical compositions may be provided to the individual by a variety of routes such as subcutaneous, topical, oral, mucosal, intravenous and intramuscular.

The vaccines of the invention comprise DNA, RNA or proteins encoded by the DNA that contain the antigenic determinants necessary to induce the formation of neutralizing antibodies in the host. Such vaccines are also safe enough to be administered without danger of clinical infection; do not have toxic side effects; can be administered by an effective route; are stable; and are compatible with vaccine carriers.

The vaccines may be administered by a variety of routes, such as orally, parenterally, subcutaneously, mucosally, intravenously or intramuscularly. The dosage administered may vary with the condition, sex, weight, and age of the individual; the route of administration; and the type PV of the vaccine. The vaccine may be used in dosage forms such as capsules, suspensions, elixirs, or liquid solutions. The vaccine may be formulated with an immunologically acceptable carrier.

The vaccines are administered in therapeutically effective amounts, that is, in amounts sufficient to generate a immunologically protective response. The therapeutically effective amount may vary

- 5 -

according to the type of PV. The vaccine may be administered in single or multiple doses.

The DNA and DNA derivatives of the present invention may be used in the formulation of immunogenic compositions. Such
5 compositions, when introduced into a suitable host, are capable of inducing an immune response in the host.

The DNA or its derivatives may be used to generate antibodies. The term "antibody" as used herein includes both polyclonal and monoclonal antibodies, as well as fragments thereof, such as, Fv, Fab
10 and F(ab)2 fragments that are capable of binding antigen or hapten.

The DNA and DNA derivatives of the present invention may be used to serotype HPV infection and HPV screening. The DNA, recombinant proteins, VLP and antibodies lend themselves to the formulation of kits suitable for the detection and serotyping of HPV.
15 Such a kit would comprise a compartmentalized carrier suitable to hold in close confinement at least one container. The carrier would further comprise reagents such as HPV18 DNA, recombinant HPV protein or VLP or anti-HPV antibodies suitable for detecting a variety of HPV types. The carrier may also contain means for detection such as labeled
20 antigen or enzyme substrates or the like.

The DNA and derived proteins therefrom are also useful as molecular weight and molecular size markers.

Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore,
25 the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the HPV18 sequence but will be capable of hybridizing to HPV18 DNA even in the presence of DNA oligonucleotides with mismatches under appropriate conditions. Under alternate conditions, the
30 mismatched DNA oligonucleotides may still hybridize to the HPV18 DNA to permit identification and isolation of HPV18 encoding DNA.

The purified HPV18 DNA of the invention or fragments thereof may be used to isolate and purify homologues and fragments of HPV18 from other sources. To accomplish this, the first HPV18

- 6 -

DNA may be mixed with a sample containing DNA encoding homologues of HPV18 under appropriate hybridization conditions. The hybridized DNA complex may be isolated and the DNA encoding the homologous DNA may be purified therefrom.

5 It is known that there is a substantial amount of redundancy in the various codons which code for specific amino acids. Therefore, this invention is also directed to those DNA sequences which contain alternative codons which code for the eventual translation of the identical amino acid. For purposes of this
10 specification, a sequence bearing one or more replaced codons will be defined as a degenerate variation. Also included within the scope of this invention are mutations either in the DNA sequence or the translated protein which do not substantially alter the ultimate physical properties of the expressed protein. For example, substitution of
15 valine for leucine, arginine for lysine, or asparagine for glutamine may not cause a change in functionality of the polypeptide.

 It is known that DNA sequences coding for a peptide may be altered so as to code for a peptide having properties that are different than those of the naturally-occurring peptide. Methods of altering the DNA
20 sequences include, but are not limited to site-directed mutagenesis.

 As used herein, a "functional derivative" of HPV18 is a compound that possesses a biological activity (either functional or structural) that is substantially similar to the biological activity of HPV18. The term "functional derivatives" is intended to include the
25 "fragments," "variants," "degenerate variants," "analogs" and "homologues" or to "chemical derivatives" of HPV18. The term "fragment" is meant to refer to any polypeptide subset of HPV18. The term "variant" is meant to refer to a molecule substantially similar in structure and function to either the entire HPV18 molecule or to a
30 fragment thereof. A molecule is "substantially similar" to HPV18 if both molecules have substantially similar structures or if both molecules possess similar biological activity. Therefore, if the two molecules possess substantially similar activity, they are considered to be variants

- 7 -

even if the structure of one of the molecules is not found in the other or even if the two amino acid sequences are not identical.

The term "analog" refers to a molecule substantially similar in function to either the entire HPV18 molecule or to a
5 fragment thereof.

A variety of procedures may be used to molecularly clone HPV18 DNA. These methods include, but are not limited to, direct functional expression of the HPV18 genes following the construction of a HPV18-containing cDNA or genomic DNA library in an
10 appropriate expression vector system. Another method is to screen HPV18-containing cDNA or genomic DNA library constructed in a bacteriophage or plasmid shuttle vector with a labeled oligonucleotide probe designed from the amino acid sequence of the HPV18. An additional method consists of screening a HPV18-containing cDNA or
15 genomic DNA library constructed in a bacteriophage or plasmid shuttle vector with a partial DNA encoding the HPV18. This partial DNA is obtained by the specific polymerase chain reaction (PCR) amplification of HPV18 DNA fragments through the design of degenerate oligonucleotide primers from the amino acid sequence of
20 purified HPV18. Another method is to isolate RNA from HPV18-producing cells and translate the RNA into protein via an *in vitro* or an *in vivo* translation system. The translation of the RNA into a peptide or a protein will result in the production of at least a portion of HPV18 protein which can be identified by, for example, the activity of HPV18
25 protein or by immunological reactivity with an anti-HPV18 antibody. In this method, pools of RNA isolated from HPV18-producing cells can be analyzed for the presence of an RNA which encodes at least a portion of the HPV18. Further fractionation of the RNA pool can be done to purify the HPV18 RNA from non-HPV18 RNA. The peptide
30 or protein produced by this method may be analyzed to provide amino acid sequences which in turn are used to provide primers for production of HPV18 cDNA, or the RNA used for translation can be analyzed to provide nucleotide sequences encoding HPV18 and produce probes for the screening of a HPV18 cDNA library. These

- 8 -

methods are known in the art and can be found in, for example, Sambrook, J., Fritsch, E.F., Maniatis, T. in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 1989.

5 It is apparent that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating HPV18-encoding DNA. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells or cell lines containing HPV type 18 and genomic DNA libraries.

10 Preparation of cDNA libraries can be performed by a variety of techniques. cDNA library construction techniques can be found in Sambrook, J., *et al.*, *supra*. It is apparent that DNA encoding HPV18 may also be isolated from a suitable genomic DNA library. Construction of genomic DNA libraries can be performed by a variety
15 of techniques. Genomic DNA library construction techniques can be found in Sambrook, J., *et al.* *supra*.

 The cloned HPV18 DNA or fragments thereof obtained through the methods described herein may be recombinantly expressed by molecular cloning into an expression vector containing a
20 suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce recombinant HPV18. Techniques for such manipulations are fully described in Sambrook, J., *et al.*, *supra*, and are known in the art.

 Expression vectors are defined herein as DNA sequences
25 that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria, bluegreen algae, plant cells, insect cells, fungal cells and animal cells. Specifically designed vectors allow the shuttling of
30 DNA between hosts such as bacteria-yeast or bacteria-animal cells or bacteria-fungal cells or bacteria-invertebrate cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy

- 9 -

number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

A variety of mammalian expression vectors may be used to express HPV18 DNA or fragments thereof in mammalian cells. Commercially available mammalian expression vectors which may be suitable for recombinant HPV18 expression, include but are not limited to, pcDNA3 (Invitrogen), pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593) pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), and λ ZD35 (ATCC 37565).

A variety of bacterial expression vectors may be used to express HPV18 DNA or fragments thereof in bacterial cells. Commercially available bacterial expression vectors which may be suitable include, but are not limited to pET11a (Novagen), lambda gt11 (Invitrogen), pcDNAII (Invitrogen), pKK223-3 (Pharmacia).

A variety of fungal cell expression vectors may be used to express HPV18 or fragments thereof in fungal cells. Commercially available fungal cell expression vectors which may be suitable include but are not limited to pYES2 (Invitrogen), *Pichia* expression vector (Invitrogen), and *Hansenula* expression (Rhein Biotech, Dusseldorf, Germany).

A variety of insect cell expression vectors may be used to express HPV18 DNA or fragments thereof in insect cells. Commercially available insect cell expression vectors which may be suitable include but are not limited to pBlue Bac III (Invitrogen) and pAcUW51 (PharMingen, Inc.).

An expression vector containing DNA encoding HPV18 or fragments thereof may be used for expression of HPV18 proteins or

- 10 -

fragments of HPV18 proteins in a cell, tissues, organs, or animals (including humans). Host cells may be prokaryotic or eukaryotic, including but not limited to bacteria such as *E. coli*, fungal cells such as yeast, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to *Drosophila* and silkworm derived cell lines. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, L cells L-M(TK-) (ATCC CCL 1.3), L cells L-M (ATCC CCL 1.2), 293 (ATCC CRL 1573), Raji (ATCC CCL 86), CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C127I (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, lipofection, protoplast fusion, and electroporation. The expression vector-containing cells are clonally propagated and individually analyzed to determine whether they produce HPV18 protein. Identification of HPV18 expressing host cell clones may be done by several means, including but not limited to immunological reactivity with anti-HPV18 antibodies, and the presence of host cell-associated HPV18 activity, such as HPV18-specific ligand binding or signal transduction defined as a response mediated by the interaction of HPV18-specific ligands with the expressed HPV18 proteins.

Expression of HPV DNA fragments may also be performed using *in vitro* produced synthetic mRNA or native mRNA. Synthetic mRNA or mRNA isolated from HPV18 producing cells can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being preferred.

- 11 -

Following expression of HPV18 protein(s) in a host cell, HPV18 protein may be recovered to provide HPV18 in purified form. Several HPV18 purification procedures are available and suitable for use. As described herein, recombinant HPV18 protein may be
5 purified from cell lysates and extracts by various combinations of, or individual application of salt fractionation, ion exchange chromatography, size exclusion chromatography, hydroxylapatite adsorption chromatography and hydrophobic interaction chromatography.

10 In addition, recombinant HPV18 may be separated from other cellular proteins by use of an immunoaffinity column made with monoclonal or polyclonal antibodies specific for full length nascent HPV18, or polypeptide fragments of HPV18. Monoclonal and polyclonal antibodies may be prepared according to a variety of
15 methods known in the art. Monoclonal or monospecific antibody as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for HPV18. Homogenous binding as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope.

20 It is apparent that the methods for producing monospecific antibodies may be utilized to produce antibodies specific for HPV18 polypeptide fragments, or full-length nascent HPV18 polypeptide. Specifically, it is apparent that monospecific antibodies may be generated which are specific for the fully functional HPV18 or
25 fragments thereof.

The present invention is also directed toward methods for screening for compounds which modulate the expression of DNA or RNA encoding HPV18 as well as the function(s) of HPV18 protein(s) *in vivo*. Compounds which modulate these activities may be DNA,
30 RNA, peptides, proteins, or non-proteinaceous organic molecules. Compounds may modulate by increasing or attenuating the expression of DNA or RNA encoding HPV18, or the function of HPV18 protein. Compounds that modulate the expression of DNA or RNA encoding HPV18 or the function of HPV18 protein may be detected by a variety

- 12 -

of assays. The assay may be a simple "yes/no" assay to determine whether there is a change in expression or function. The assay may be made quantitative by comparing the expression or function of a test sample with the levels of expression or function in a standard sample.

5 Kits containing HPV18 DNA, fragments of HPV18 DNA, antibodies to HPV18 DNA or HPV18 protein, HPV18 RNA or HPV18 protein may be prepared. Such kits are used to detect DNA which hybridizes to HPV18 DNA or to detect the presence of HPV18 protein(s) or peptide fragments in a sample. Such characterization is
10 useful for a variety of purposes including but not limited to forensic analyses and epidemiological studies.

 Nucleotide sequences that are complementary to the HPV18 encoding DNA sequence may be synthesized for antisense therapy. These antisense molecules may be DNA, stable derivatives
15 of DNA such as phosphorothioates or methylphosphonates, RNA, stable derivatives of RNA such as 2'-O-alkylRNA, or other HPV18 antisense oligonucleotide mimetics. HPV18 antisense molecules may be introduced into cells by microinjection, liposome encapsulation or by expression from vectors harboring the antisense sequence. HPV18
20 antisense therapy may be particularly useful for the treatment of diseases where it is beneficial to reduce HPV18 activity.

 The term "chemical derivative" describes a molecule that contains additional chemical moieties which are not normally a part of the base molecule. Such moieties may improve the solubility, half-
25 life, absorption, etc. of the base molecule. Alternatively the moieties may attenuate undesirable side effects of the base molecule or decrease the toxicity of the base molecule. Examples of such moieties are described in a variety of texts, such as Remington's Pharmaceutical Sciences.

30 Compounds identified according to the methods disclosed herein may be used alone at appropriate dosages defined by routine testing in order to obtain optimal inhibition of the HPV18 or its activity while minimizing any potential toxicity. In addition, co-administration or sequential administration of other agents may be desirable.

- 13 -

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in several divided doses. Furthermore, compounds for the present invention may be administered via a
5 variety of routes including but not limited to intranasally, orally, transdermally or by suppository.

For combination treatment with more than one active agent, where the active agents are in separate dosage formulations, the active agents can be administered concurrently, or they each can be
10 administered at separately staggered times.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of
15 administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of drug within the
20 range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

In the methods of the present invention, the compounds
25 herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrup,
30 suppositories, gels and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as

- 14 -

ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

For liquid forms the active drug component can be combined in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. Other dispersing agents which may be employed include glycerin and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Topical preparations containing the active drug component can be admixed with a variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the

- 15 -

present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethylenoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

The following examples illustrate the present invention without, however, limiting the same thereto.

15 EXAMPLE 1

Cloning of HPV18 genome

Total genomic DNA was prepared from the human cervical carcinoma-derived cell line, SW756 (Freedman, R. S., *et al.*, 1982, *In Vitro*, Vol 18, pages 719-726) by standard techniques. The DNA was digested with EcoR1 and electrophoresed through a 0.8% low-melting temperature, agarose preparative gel. A gel slice was excised corresponding to DNA fragments approximately 12 kbp in length. The agarose was digested using AgaraseTM enzyme (Boehringer Mannheim, Inc.) and the size-fractionated DNA was precipitated, dephosphorylated and ligated with EcoR1 digested lambda EMBL4 arms (Stratagene, Inc.). The lambda library was packaged using Gigapack II Gold packaging extract (Stratagene, Inc.). HPV18-positive clones were identified using a 700 bp, HPV18 L1 DNA probe that was generated by polymerase chain reaction (PCR) using SW756 DNA as template and oligonucleotide primers that were designed based on the published HPV18 L1 DNA sequence (Cole and Danos, 1987, *J. Mol. Biol.*, Vol. 193:599-608; Genbank Accession #X05015). A HPV18-positive, lambda clone was isolated that contained a 12 kbp EcoR1 fragment insert and was designated as #187-1.

EXAMPLE 2Construction of Yeast Expression Vectors

The HPV18 L1 open reading frame (ORF) was amplified by
 5 PCR using clone #187-1 as template, Vent polymeraseTM (New England
 Biolabs, Inc.), 10 cycles of amplification (94°C, 1 min; 50°C, 1 min;
 72°C, 2 min) and the following oligonucleotide primers which contain
 flanking BglII sites (underlined):

sense primer,

10 5'-GAAGATCTCACAAAACAAAATGGCTTTGTGG
 CGGCCTAGTG-3',

antisense primer, 5'-GAAGATCTTTACTTCCTGGCACGTAC
 ACGCACACGC-3'.

The sense primer introduces a yeast non-translated leader sequence
 15 (Kniskern, *et al.*, 1986, *Gene*, Vol. 46:135-141) immediately upstream to
 the HPV18 L1 initiating methionine codon (highlighted in bold print).
 The 1.5 kbp L1 PCR product was digested with BglII and gel purified.

The pGAL1-10 yeast expression vector was constructed by
 isolating a 1.4 kbp SphI fragment from a pUC18/bidirectional *GAL*
 20 promoter plasmid which contains the *Saccharomyces cerevisiae* divergent
GAL1-GAL10 promoters from the plasmid pBM272 (provided by Mark
 Johnston, Washington University, St. Louis, MO). The divergent
 promoters are flanked on each side by a copy of the yeast *ADHI*
 transcriptional terminator, a BamHI cloning site located between the
 25 *GAL1* promoter and the first copy of the *ADHI* transcriptional terminator
 and a SmaI cloning site located between the *GAL10* promoter and the
 second copy of the *ADHI* transcriptional terminator. A yeast shuttle
 vector consisting of pBR322, the yeast *LEU2d* gene, and the yeast *2u*
 plasmid (gift of Benjamin Hall, University of Washington, Seattle, WA)
 30 was digested with SphI and ligated with the 1.4 kbp SphI divergent *GAL*
 promoter fragment resulting in pGAL1-10. pGAL1-10 was linearized
 with BamHI which cuts between the *GAL1* promoter and the *ADHI*
 transcription terminator. The BamHI digested vector and the BglII
 digested HPV18 L1 PCR fragment were ligated and used to transform

- 17 -

E. coli DH5 cells (Gibco BRL, Inc.). A pGAL1-10 plasmid was isolated which contains the HPV18 L1 gene and was designated p191-6.

A yeast expression vector that co-expresses both the HPV18 L1 and L2 genes was constructed. Plasmid p191-6 (pGAL1-10 + HPV18 L1) was digested with SmaI which cuts between the *GAL10* promoter and the *ADHI* transcription terminator. The 1.4 kbp HPV18 L2 gene was amplified by PCR as described above using the following oligonucleotide primers which contain flanking SmaI sites (underlined):
sense primer,

10 5'-TCCCCCGGGCACAACAAAATG
 GTATCCCACCGTGCCGCACGAC-3',

antisense primer,

5'-TCCCCCGGGCTAGGCCGCCACAAAGCCATCTGC-3'.

The sense primer introduces a yeast non-translated leader sequence (Kniskern *et al.*, 1986, *supra*) immediately upstream to the HPV18 L2 initiating methionine codon (highlighted in bold print). The PCR fragment was digested with SmaI, gel purified and ligated with the SmaI digested p191-6 plasmid. A pGAL1-10 plasmid containing both the HPV18 L1 and L2 genes was isolated and designated, p195-11.

20

EXAMPLE 3

Typing of Clinical Samples

Cervical biopsy samples were collected at the Veterans Administration Medical Center in Indianapolis, IN (courtesy of Dr. Darron Brown) and at the Albert Einstein Medical Center in Philadelphia, PA (courtesy of Dr. Joan Adler) and were frozen at - 20°C. DNA was isolated as described by Brown *et al.*, 1993 (Brown, D. *et al.*, 1993, *J. Clin. Microbiol.*, Vol. 31:2667-2673). Briefly, clinical specimens were processed with a Braun mikro-dismembrator II (B. Braun Instruments, Melsungen, Germany) and solubilized in buffer containing 10 mM EDTA and 0.6% (w/v) sodium dodecyl sulfate (SDS). Samples were adjusted to 20 mM Tris pH 7.4 and protein was digested with 50 mcg/mL Proteinase K in the presence of 0.1 mcg/mL RNase A followed by extraction with

25

30

- 18 -

phenol/chloroform/isoamyl alcohol. DNA was ethanol precipitated and quantified by spectrophotometry.

The DNA samples were screened for the presence of HPV18 by PCR and Southern blot analyses. A 256 bp segment of the HPV18 L1 ORF was amplified by PCR using the following oligonucleotide primers:

5'-CAATCCTTATATATTAAAGGCACAGGTATG-3',

antisense primer,

5'-CATCATATTGCCAGGTACAGGAGACTGTG-3'.

The PCR conditions were according to the manufacturer's recommendations for the AmpliTaqTM DNA Polymerase/GeneAmpTM kit (Perkin Elmer Corp.) except that 0.5 µl of clinical sample DNA was used as template and 10 pmoles of each primer, 2 mM dNTPs and 2.0 mM MgCl₂ were in the final reaction mixture. A 2 min, 94°C denaturation step was followed by 40 cycles of amplification (94°C, 1 min; 45°C, 1 min; 72°C, 1 min).

The PCR products were electrophoresed through a 3.0% agarose gel, blotted onto nylon membranes and hybridized with a ³²P-labeled HPV18 L1-specific oligonucleotide probe.

EXAMPLE 4

DNA Sequencing of L1 and L2 genes

The HPV18 L1 and L2 genes in clones #187-1, p191-6 and p195-11 were sequenced using the PRIZM Sequencing kit and the automated DNA ABI Sequencer #373A (Applied Biosystems). To obtain a consensus HPV18 sequence, portions of the L1 gene DNA were amplified by PCR from human clinical isolates, sequenced and compared to the claimed and published sequences. A 256 bp fragment (nucleotides 817-1072) was amplified from each clinical DNA isolate for this purpose using the oligonucleotides and heating cycles described in Example 3.

The following primers,

5'-GAAGATCTCACAAAACAAAATGGCTTTGTGGCGGCCTAGTG-3' and 5'-CCTAACGTCCTCAGAAACATTAGAC-3' were used to

amplify an amino-terminal 432 bp portion of L1 DNA (nucleotides 1-

- 19 -

431) using the heating cycles described in Example 3. Both PCR products were ligated separately with plasmid pCRII (Invitrogen Corp.) using the reagents and procedures recommended by the manufacturer. Plasmid DNA was isolated from the transformants and those containing
5 EcoRI inserts were sequenced.

EXAMPLE 5

Analysis of DNA and Deduced Amino Acid Sequences

10 The nucleotide and deduced amino acid (aa) sequences of the claimed HPV18 L1 are shown in Figure 1. The DNA sequence was derived from a consensus of clones #187-1, p191-6 and p195-11. A comparison of the claimed HPV18 L1 nucleotide sequence with the published HPV18 L1 sequence (Genbank Accession #X05015) identified
15 20 bp changes out of 1524 bps. Five of the nucleotide changes (C to G at position 89, C to A at 263, C to G at 848, G to A at 967 and C to G at 1013) result in amino acid substitutions. The five residue differences from published are P to R at aa positions 30, 283 and 338, T to N at aa 88 and V to I at aa 323 (Figure 2). Positions 88 and 323 represent
20 conservative changes while the three P to R changes may substantially alter the physical properties of the expressed L1 protein.

A comparison of the amino acid sequences derived from clinical isolates (numbers 354, 556, 755, 697, 795 and 23) with the claimed sequence and the published sequence is shown in Figure 2.
25 There are four locations where the clinical isolates and the claimed sequence differ from the published sequence. Positions 30, 283 and 338 encode arginine (R) in all the isolates found to date, including the claimed sequence. This is in sharp contrast to the published sequence which has prolines (P) at each of these locations. Furthermore, position 88 is an
30 asparagine (N) in the isolates and the claimed sequence but is a threonine (T) in the published sequence. The last difference, position 323, was found to be a valine (V) in many of the clinical isolates and the published strain versus an isoleucine (I) in the claimed sequence and one of the isolates (#23). The conclusion is that the claimed sequence reflects the
35 predominant viral sequences that are associated with clinical infections

- 20 -

and the absence of isolates containing any of the position 30, 283 or 338 prolines of the published sequence suggests that the published clone is either an artefact or an inconsequential subtype.

The nucleotide and deduced aa sequences of HPV18 L2 were derived from a consensus sequence of clones #187-1 and p195-11 and are shown in Figure 3. A comparison of the L2 nucleotide sequence with the published HPV18 sequence (Genbank Accession #X05015) identified 40 bp changes out of 1389 bps. The bp differences result in 14 changes at the aa level: P to S at aa 29, P to N at aa 33, A to S at aa 177, D to E at aa 266, D to N at aa 270, D to G at aa 346, M to I at 355, V to M at aa 359, S to P at aa 365, F to S at aa 369, F to V at aa 371, F to S at aa 372, K to T at aa 373 and S to P at aa 409.

EXAMPLE 6

Generation of HPV18 L2 Antiserum

HPV18 L2 specific antibodies were prepared in goats using a *trpE*-HPV18 L2 fusion protein expressed in *E. coli*. The full-length L2 ORF was amplified by PCR using oligonucleotide primers providing HindIII and BamHI sites flanking the 5'- and 3'- ends, respectively. The L2 fragment was inserted into the HindIII-BamHI digested, pATH23 expression plasmid (Koerner *et al.*, 1991, *Meth. Enzymol.* Vol. 194:477-490). The fusion protein was expressed in *E. coli* RR1 cells (Gibco BRL, Inc.) after induction with 3-b-indoleacrylic acid. The insoluble fraction was analyzed by SDS-PAGE followed by staining with Coomassie Blue. The *trpE*-L2 fusion protein accounted for the major portion of the *E. coli* insoluble fraction. Goats were immunized with the *trpE*-L2 fusion protein according to the standard protocol of Pocono Rabbit Farm and Laboratory, Inc. for fusion protein antigens (Protein Rabbit Farm, Canadensis, PA).

- 21 -

EXAMPLE 7

Preparation of Yeast Strain U9

Saccharomyces cerevisiae strain 2150-2-3 (*MAT α* , *leu2-04*, *ade1*, *cir*^o) was obtained from Dr. Leland Hartwell (University of Washington, Seattle, WA). Cells of strain 2150-2-3 were propagated overnight at 30°C in 5 mL of YEHD medium (Carty *et al.*, *J. Ind Micro* 2 (1987) 117-121). The cells were washed 3 times in sterile, distilled water, resuspended in 2 mL of sterile distilled water, and 0.1 mL of cell suspension was plated onto each of six 5-fluoro-orotic acid (FOA) plates in order to select for *ura3* mutants (Cold Spring Harbor Laboratory Manual for Yeast Genetics). The plates were incubated at 30°C. The medium contained per 250 mL distilled water: 3.5 g Difco Yeast Nitrogen Base without amino acids and ammonium sulfate; 0.5 g 5-Fluoro-orotic acid; 25 mg Uracil; and 10.0 g Dextrose.

The medium was sterilized by filtration through 0.2 μ m membranes and then mixed with 250 mL of 4% Bacto-Agar (Difco) maintained at 50°C, 10 mL of a 1.2 mg/mL solution of adenine, and 5 mL of L-leucine solution (180 mg/ 50 mL). The resulting medium was dispensed at 20 mL per petri dish.

After 5 days of incubation, numerous colonies had appeared. Single colonies were isolated by restreaking colonies from the initial FOA plates onto fresh FOA plates which were then incubated at 30°C. A number of colonies from the second set of FOA plates were tested for the presence of the *ura3* mutation by replica-plating onto both YEHD plates and uracil-minus plates. The desired result was good growth on YEHD and no growth on uracil-minus medium. One isolate (U9) was obtained which showed these properties. It was stored as a frozen glycerol stock (strain #325) at -70°C for later use.

EXAMPLE 8

Preparation of a Vector for disruption of the Yeast *MNN9* gene

In order to prepare a vector for disruption of the *MNN9* gene, it was necessary to first clone the *MNN9* gene from *S. cerevisiae* genomic

- 22 -

DNA. This was accomplished by standard Polymerase Chain Reaction (PCR) technology. A 5' sense primer and 3' antisense primer for PCR of the full-length *MNN9* coding sequence were designed based on the published sequence for the yeast *MNN9* gene (Zymogenetics: EPO Patent Application No. 88117834.7, Publication No. 0-314-096-A2). The following oligodeoxynucleotide primers containing flanking HindIII sites (underlined) were used:

10 sense primer: 5'-CTT AAA GCT TAT GTC ACT TTC TCT TGT ATC
G-3'

antisense primer: 5'-TGA TAA GCT TGC TCA ATG GTT CTC TTC
CTC-3'.

15 The initiating methionine codon for the *MNN9* gene is highlighted in bold print. The PCR was conducted using genomic DNA from *S. cerevisiae* strain JRY188 as template, Taq DNA polymerase (Perkin Elmer) and 25 cycles of amplification (94°C 1 min., 37°C 2 min., 72°C 3 min.). The resulting 1.2 kbp PCR fragment was digested with
20 HindIII, gel-purified, and ligated with HindIII-digested, alkaline-phosphatase treated pUC13 (Pharmacia). The resulting plasmid was designated p1183.

In order to disrupt the *MNN9* gene with the yeast *URA3* gene, the plasmid pBR322-*URA3* (which contains the 1.1 Kbp HindIII fragment encoding the *S. cerevisiae URA3* gene subcloned into the
25 HindIII site of pBR322) was digested with HindIII and the 1.1 kbp DNA fragment bearing the functional *URA3* gene was gel-purified, made blunt-ended with T4 DNA polymerase, and then ligated with PmlI-digested plasmid p1183 (PmlI cuts within the *MNN9* coding sequence).
30 The resulting plasmid p1199 contains a disruption of the *MNN9* gene by the functional *URA3* gene.

- 23 -

EXAMPLE 9

Construction of U9-derivative strain 1372 containing disruption of *MNN9* gene

5 For disruption of the *MNN9* gene in strain U9 (#325), 30 µg of plasmid p1199 were digested with HindIII to create a linear *mn9::URA3* disruption cassette. Cells of strain 325 were transformed with the HindIII-digested p1199 DNA by the spheroplast method (Hinnen *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75:1929-1933) and transformants
10 were selected on a synthetic agar medium lacking uracil and containing 1.0 M sorbitol. The synthetic medium contained, per liter of distilled water: Agar, 20 g; Yeast nitrogen base w/o amino acids, 6.7 g; Adenine, 0.04 g; L-tyrosine, 0.05 g; Sorbitol, 182 g; Glucose, 20 g; and Leucine Minus Solution #2, 10 ml. Leucine Minus Solution #2 contains per liter
15 of distilled water: L-arginine, 2 g; L-histidine, 1 g; L-Leucine, 6 g; L-Isoleucine, 6 g; L-lysine, 4 g; L-methionine, 1 g; L-phenylalanine, 6 g; L-threonine, 6 g; L-tryptophan, 4 g.

The plates were incubated at 30°C for five days at which time numerous colonies had appeared. Chromosomal DNA preparations
20 were made from 10 colonies and then digested with EcoRI plus HindIII. The DNA digests were then evaluated by Southern blots (J. Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd edition, Cold Spring Harbor Laboratory Press, 1989) using the 1.2 kbp HindIII fragment bearing the *MNN9* gene (isolated from plasmid p1199) as a
25 probe. An isolate was identified (strain #1372) which showed the expected DNA band shifts on the Southern blot as well as the extreme clumpiness typically shown by *mn9* mutants.

EXAMPLE 10

Construction of a Vector for Disruption of Yeast *HIS3* Gene

30 In order to construct a disruption cassette in which the *S. cerevisiae* *HIS3* gene is disrupted by the *URA3* gene, the plasmid YEp6 (K. Struhl *et al.*, 1979, *Proc. Natl. Acad. Sci., USA* 76:1035) was digested
35 with BamHI and the 1.7 kbp BamHI fragment bearing the *HIS3* gene was

- 24 -

gel-purified, made blunt-ended with T4 DNA polymerase, and ligated with pUC18 which had been previously digested with BamHI and treated with T4 DNA polymerase. The resulting plasmid (designated p1501 or pUC18-HIS3) was digested with NheI (which cuts in the *HIS3* coding sequence), and the vector fragment was gel-purified, made blunt-ended with T4 DNA polymerase, and then treated with calf intestine alkaline phosphatase. The *URA3* gene was isolated from the plasmid pBR322-*URA3* by digestion with HindIII and the 1.1 kbp fragment bearing the *URA3* gene was gel-purified, made blunt-ended with T4 DNA polymerase, and ligated with the above pUC18-HIS3 NheI fragment. The resulting plasmid (designated pUC18-his3::*URA3* or p1505) contains a disruption cassette in which the yeast *HIS3* gene is disrupted by the functional *URA3* gene.

EXAMPLE 11

Construction of Vector for Disruption of Yeast *PRB1* Gene by the *HIS3* Gene

Plasmid FP8 Δ H bearing the *S. cerevisiae PRB1* gene was provided by Dr. E. Jones of Carnegie-Mellon Univ. (C. M. Moehle *et al.*, 1987, *Genetics* 115:255-263). It was digested with HindIII plus XhoI and the 3.2 kbp DNA fragment bearing the *PRB1* gene was gel-purified and made blunt-ended by treatment with T4 DNA polymerase. The plasmid pUC18 was digested with BamHI, gel-purified and made blunt-ended by treatment with T4 DNA polymerase. The resulting vector fragment was ligated with the above *PRB1* gene fragment to yield the plasmid pUC18-*PRB1*. Plasmid YEp6, which contains the *HIS3* gene, was digested with BamHI. The resulting 1.7 kbp BamHI fragment bearing the functional *HIS3* gene was gel-purified and then made blunt-ended by treatment with T4 DNA polymerase. Plasmid pUC18-*PRB1* was digested with EcoRV plus NcoI which cut within the *PRB1* coding sequence and removes the protease B active site and flanking sequence. The 5.7 kbp EcoRV-NcoI fragment bearing the residual 5' and 3'-portions of the *PRB1* coding sequence in pUC18 was gel-purified, made blunt-ended by treatment with T4 DNA polymerase, dephosphorylated with calf intestine alkaline

- 25 -

phosphatase, and ligated with the blunt-ended *HIS3* fragment described above. The resulting plasmid (designated pUC18-prb1::HIS3, stock #1245) contains the functional *HIS3* gene in place of the portion of the *PRB1* gene which had been deleted above.

5

EXAMPLE 12

Construction of a U9-related Yeast Strain containing disruptions of both the *MNN9* and *PRB1* Genes

10 The U9-related strain 1372 which contains a *MNN9* gene disruption was described in Example 9. Clonal isolates of strain 1372 were passaged on FOA plates (as described in Example 7) to select *ura3* mutants. A number of *ura3* isolates of strain 1372 were obtained and one particular isolate (strain 12930-190-S1-1) was selected for subsequent
15 disruption of the *HIS3* gene. The pUC18-his3::URA3 gene disruption vector (p1505) was digested with XbaI plus EcoRI to generate a linear *his3::URA3* disruption cassette and used for transformation of strain 12930-190-S1-1 by the lithium acetate method [Methods in Enzymology, 194:290 (1991)]. Ura⁺ transformants were selected on synthetic agar
20 medium lacking uracil, restreaked for clonal isolates on the same medium, and then replica-plated onto medium lacking either uracil or histidine to screen for those isolates that were both Ura⁺ and His⁻. One isolate (strain 12930-230-1) was selected for subsequent disruption of the *PRB1* gene. The *PRB1* gene disruption vector (pUC18-prb1::HIS3, stock
25 #1245) was digested with SacI plus XbaI to generate a linear *prb1::HIS3* disruption cassette and used for transformation of strain 12930-230-1 by the lithium acetate method. His⁺ transformants were selected on agar medium lacking histidine and restreaked on the same medium for clonal isolates. Genomic DNA was prepared from a number of the resulting
30 His⁺ isolates, digested with EcoRI, and then electrophoresed on 0.8% agarose gels. Southern blot analyses were then performed using a radio-labeled 617 bp probe for the *PRB1* gene which had been prepared by PCR using the following oligodeoxynucleotide primers:

35 5' TGG TCA TCC CAA ATC TTG AAA 3'
 5' CAC CGT AGT GTT TGG AAG CGA 3'

- 26 -

Eleven isolates were obtained which showed the expected hybridization of the probe with a 2.44 kbp *prb1::HIS3* DNA fragment. This was in contrast to hybridization of the probe with the 1.59 kbp fragment for the wild-type *PRB1* gene. One of these isolates containing the desired *prb1::HIS3* disruption was selected for further use and was designated strain #1558.

EXAMPLE 13

10 Expression of HPV18 L1 and L2 in Yeast

Plasmids p191-6 (pGAL1-10 + HPV18 L1) and p195-11 (pGAL1-10 + HPV18 L1 + L2) were used to transform *S. cerevisiae* strain #1558 (*MATa*, *leu2-04*, *prb1::HIS3*, *mnn9::URA3*, *ade1*, *cir⁰*). Clonal isolates were grown at 30°C in YEHD medium containing 2% galactose for 88 hours. After harvesting the cells, the cell pellets were broken with glass beads and cell lysates analyzed for the expression of HPV18 L1 and/or HPV18 L2 protein by immunoblot analysis. Samples containing 25 ug of total cellular protein were electrophoresed through 10% Tris-Glycine gels (Novex, Inc.) under denaturing conditions and electroblotted onto nitrocellulose filters. L1 protein was immunodetected using rabbit antiserum raised against a *trpE*-HPV11 L1 fusion protein as primary antibody (Brown *et al.*, 1994, *Virology* 201:46-54) and horseradish peroxidase (HRP)-linked donkey anti-rabbit IgG (Amersham, Inc.) as secondary antibody. The filters were processed using the chemiluminescent ECLTM Detection Kit (Amersham, Inc.). A 50-55 KDa L1 protein band was detected in both the L1 and L1 + L2 coexpressor yeast clones (strains 1725 and 1727, respectively) and not in the negative control (pGAL1-10 without L1 or L2 genes) (Figure 4).

The HPV18 L2 protein was detected by Western analysis using goat polyclonal antiserum raised against a *trpE*-HPV18 L2 fusion protein as primary antibody followed by HRP-conjugated, rabbit anti-goat IgG (Kirkegaard and Perry Laboratories, Gaithersburg, MD). The filters were processed as described above. The L2 protein was detected as a 75 kDa protein band in the L1 + L2 coexpressor yeast clone (strain

- 27 -

1727) but not in either the negative control or the L1 expressor clone (Figure 5).

EXAMPLE 14

5

Fermentation of HPV 18 L1 (strain 1725) and 18 L1+ΔL2 (strain 1727).

Surface growth of a plate culture of strains 1725 and 1727 was aseptically transferred to a leucine-free liquid medium containing (per L): 8.5 g Difco yeast nitrogen base without amino acids and ammonium sulfate; 0.2 g adenine; 0.2 g uracil; 10 g succinic acid; 5 g ammonium sulfate; 40 g glucose; 0.25 g L-tyrosine; 0.1 g L-arginine; 0.3 g L-isoleucine; 0.05 g L-methionine; 0.2 g L-tryptophan; 0.05 g L-histidine; 0.2 g L-lysine; 0.3 g L-phenylalanine; this medium was adjusted to pH 5.0-5.3 with NaOH prior to sterilization. After growth at 28°C, 250 rpm on a rotary shaker, frozen culture vials were prepared by adding sterile glycerol to a final concentration of 17% (w/v) prior to storage at - 70°C (1 mL per cryovial). Inocula were developed in the same medium (500 mL per 2-L flask) and were started by transferring the thawed contents of a frozen culture vial and incubating at 28°C, 250 rpm on a rotary shaker for 29 hr. Fermentations of each strain used a New Brunswick SF-116 fermentor with a working volume of 10 L after inoculation. The production medium contained (per L): 20 g Difco yeast extract; 10 g Sheffield HySoy peptone; 20 g glucose; 20 g galactose; 0.3 mL Union Carbide UCON LB-625 antifoam; the medium was adjusted to pH 5.3 prior to sterilization. The entire contents (500 mL) of the 2-L inoculum flask was transferred to the fermentor which was incubated at 28°C, 5 L air per min, 400 rpm, 3.5 psi pressure. Agitation was increased as needed to maintain dissolved oxygen levels of greater than 40% of saturation. Progress of the fermentation was monitored by off-line glucose measurements (Beckman Glucose 2 Analyzer) and on-line mass spectrometry (Perkin-Elmer 1200). After incubation for 66 hr, cell densities of 9.5 to 9.7 g dry cell weight per L were reached. The cultures were concentrated by hollow fiber filtration (Amicon H5MP01-43 cartridge in an Amicon DC-10 filtration system) to ca. 2 L, diafiltered with 2 L phosphate-buffered saline, and concentrated further (to ca. 1 L)

- 28 -

before dispensing into 500-mL centrifuge bottles. Cell pellets were collected by centrifugation at 8,000 rpm (Sorval GS-3 rotor) for 20 min at 4°C. After decanting the supernatant, the pellets (total 191 to 208 g wet cells) were stored at - 70°C until use.

5

EXAMPLE 15

Purification of Recombinant HPV Type 18 L1 Capsid Proteins

All steps performed at 4°C unless noted.

10 Cells were stored frozen at -70°C. Frozen cells (wet weight = 126 g) were thawed at 20-23°C and resuspended in 70 mL "Breaking Buffer" (20 mM sodium phosphate, pH 7.2, 100 mM NaCl). The protease inhibitors PMSF and pepstatin A were added to final concentrations of 2 mM and 1.7 µM, respectively. The cell slurry was
15 broken at a pressure of approximately 16,000 psi by 4 passes in a M110-Y Microfluidizer (Microfluidics Corp., Newton, MA). The broken cell slurry was centrifuged at 12,000 x g for 40 min to remove cellular debris. The supernatant liquid containing L1 antigen was recovered.

The supernatant liquid was diluted 1:5 by addition of Buffer
20 A (20 mM MOPS, pH 7.0) and applied to an anion exchange capture column (9.0 cm ID x 3.9 cm) of Fractogel[®] EMD TMAE-650 (M) resin (EM Separations, Gibbstown, NJ) equilibrated in Buffer A. Following a wash with Buffer A, the antigen was eluted with a gradient of 0 - 1.0 M NaCl in Buffer A. Column fractions were assayed for total protein by the
25 Bradford method. Fractions were then analyzed at equal total protein loadings by Western blotting and SDS-PAGE with silver stain detection.

TMAE fractions showing comparable purity and enrichment of L1 protein were pooled. The antigen was concentrated by ammonium sulfate fractionation. The solution was adjusted to 35% saturated
30 ammonium sulfate by adding solid reagent while gently stirring over 10 min. The sample was placed on ice and precipitation allowed to proceed for 4 hours. The sample was centrifuged at 16,000 x g for 45 min. The pellet was resuspended in 20.0 mL PBS (6.25 mM Na phosphate, pH 7.2, 150 mM NaCl).

- 29 -

The resuspended pellet was chromatographed on a size exclusion column (2.6 cm ID x 89 cm) of Sephacryl 500 HR resin (Pharmacia, Piscataway, NJ). Running buffer was PBS. Fractions were analyzed by western blotting and SDS-PAGE with silver stain detection.

- 5 The purest fractions were pooled. The resulting pool was concentrated in a 50 mL stirred cell using 43 mm YM-100 flat-sheet membranes (Amicon, Beverly, MA) at a N₂ pressure of 4-6 psi.

- 10 Final product was analyzed by western blotting and SDS-PAGE with colloidal Coomassie detection. The L1 protein was estimated to be 50-60% homogeneous. The identity of L1 protein was confirmed by western blotting. The final product was aliquoted and stored at -70°C. This process resulted in a total of 12.5 mg protein.

Bradford Assay for Total Protein

- 15 Total protein was assayed using a commercially available Coomassie Plus [®] kit (Pierce, Rockford, IL). Samples were diluted to appropriate levels in Milli-Q-H₂O. Volumes required were 0.1 mL and 1.0 mL for the standard and microassay protocols, respectively. For both protocols, BSA (Pierce, Rockford, IL) was used to generate the standard curve. Assay was performed according to manufacturer's
- 20 recommendations. Standard curves were plotted using CricketGraph [®] software on a Macintosh IIfx computer.

SDS-PAGE and Western Blot Assays

- 25 All gels, buffers, and electrophoretic apparatus were obtained from Novex (San Diego, CA) and were run according to manufacturer's recommendations. Briefly, samples were diluted to equal protein concentrations in Milli-Q-H₂O and mixed 1:1 with sample incubation buffer containing 200 mM DTT. Samples were incubated 15
- 30 min at 100°C and loaded onto pre-cast 12% Tris-glycine gels. The samples were electrophoresed at 125V for 1 hr 45 min. Gels were developed using either silver staining by a variation of the method of Heukeshoven and Dernick [*Electrophoresis*, 6 (1985) 103-112] or

- 30 -

colloidal Coomassie staining using a commercially obtained kit (Integrated Separation Systems, Natick, MA).

For western blots, proteins were transferred to PVDF membranes at 25V for 40 min. Membranes were washed with Milli-Q-H₂O and air-dried. Primary antibody was polyclonal rabbit antiserum raised against a TrpE-HPV11L1 fusion protein (gift of Dr. D. Brown). Previous experiments had shown this antiserum to cross react with HPV type 18 L1 on western blots. The antibody solution was prepared by dilution of antiserum in blotting buffer (5% non-fat milk in 6.25 mM Na phosphate, pH 7.2, 150 mM NaCl, 0.02% NaN₃). Incubation was for at least 1 hour at 20-23°C. The blot was washed for 1 min each in three changes of PBS (6.25 mM Na phosphate, pH 7.2, 150 mM NaCl). Secondary antibody solution was prepared by diluting goat anti-rabbit IgG alkaline phosphatase-linked conjugate antiserum (Pierce, Rockford, IL) in blotting buffer. Incubation proceeded under the same conditions for at least 1 hour. Blots were washed as before and detected using a 1 step NBT/BCIP substrate (Pierce, Rockford, IL).

EXAMPLE 16

Electron Microscopic Studies

For EM analysis (Structure Probe, West Chester, PA), an aliquot of each sample was placed on 200-mesh carbon-coated copper grids. A drop of 2% phosphotungstic acid (PTA), pH 7.0 was placed on the grid for 20 seconds. The grids were allowed to air dry prior to transmission EM examination. All microscopy was done using a JEOL 100CX transmission electron microscope (JEOL USA, Inc.) at an accelerating voltage of 100 kV. The micrographs generated have a final magnification of 100,000x. Virus-like particles were observed in the 50-55 nm diameter size range in the yeast sample harboring the HPV18 L1 expression plasmid (Figure 6). No VLPs were observed in the yeast control samples.

EXAMPLE 17Sub-cloning of the HPV18 cDNA into expression vectors

5 The cDNA encoding HPV18 is sub-cloned into several vectors for expression of the HPV18 protein in transfected host cells and for *in vitro* transcription/translation. These vectors include pBluescript II SK+ (where expression is driven by T7 or T3 promoters) pcDNA I/Amp (where expression is driven by the cytomegalovirus (CMV) promoter), pSZ9016-1 (where expression is driven by the HIV long terminal repeat (LTR) promoter) and the baculovirus transfer vector pAcUW51 (PharMingen, Inc.) (where expression is driven by the polyhedrin (PH) promoter) for producing recombinant baculovirus containing the HPV18 encoding DNA sequence.

15 a) pBluescript II SK+:HPV18. The full length HPV18 cDNA clone is retrieved from lambda bacteriophage by limited *Eco* RI digestion and ligated into *Eco* RI-cut, CIP-treated pBluescript II SK+. Separate subclones are recovered in which the sense orientation of HPV18 followed either the T7 or T3 promoters.

20 b) pcDNA I/Amp:HPV18. To facilitate directional cloning, HPV18 is excised from a purified plasmid preparation of pBluescript II SK+:HPV18 in which the HPV18 DNA sequence is downstream of the T7 promoter using *Eco* RV and *Xba* I. The resulting *Eco* RV, *Xba* I HPV18 fragment is purified and ligated into *Eco* RV-cut, *Xba* I-cut, CIP-treated pcDNA I/Amp such that the HPV18 encoding DNA is downstream of the CMV promoter.

25 c) pSZ9016-1:HPV18. HPV18 is excised from pBluescript II SK+:HPV18 by limited *Eco* RI digestion and subsequent purification of the 1.3 Kb fragment from agarose gels. The resulting *Eco* RI HPV18 fragment is ligated into *Eco* RI-cut, CIP-treated pSZ9016-1. Subclones are selected in which the sense orientation of HPV18 is downstream of the HIV LTR promoter.

30 d) pAcUW51:HPV18 L1. The full-length HPV18 L1 ORF was amplified by PCR from clone #187-1 using oligonucleotide primers providing flanking *Bgl*II sites. The L1 gene was inserted into the BamHI site of the baculovirus transfer vector, pAcUW51 (PharMingen).

35

- 32 -

Inc.), under control of the polyhedrin promoter. Recombinant baculoviruses were generated containing the HPV18 L1 expression cassette according to the procedures described in the Pharmingen Manual. Recombinant clones were purified by limiting dilution and dot blot hybridization.

EXAMPLE 18

10 Expression Of The HPV18 Polypeptide By *In Vitro* Transcription/ Translation And By Transfection Into Host Cells

Vectors containing HPV DNA sequences are used to drive the translation of the HPV18 polypeptide in rabbit reticulocyte lysates, mammalian host cells, and in baculovirus infected insect cells. The experimental procedures are essentially those outlined in the manufacturers' instructions.

a) *In vitro* Transcription/Translation. pBluescript III SK+:HPV18 plasmid DNA (with HPV18 in the T7 orientation) is linearized by *Bam* HI digestion downstream of the HPV18 insert. The linearized plasmid is purified and used as a template for run-off transcription using T7 RNA polymerase in the presence of m7G(5')ppp(5')G. The resulting capped HPV18 transcripts are purified by LiCl precipitation and used to drive the translation of HPV18 in nuclease-pretreated rabbit reticulocyte lysate in the presence of L-[³⁵S]methionine.

b) Expression in Mammalian Cells. The HPV18 protein is expressed in mammalian host cells following transfection with either pcDNA I/Amp:HPV18 (under control of the CMV promoter) or pSZ9016-1:HPV18 (under control of the HIV LTR promoter). In the latter case (pSZ9016-1:HPV18), cells are co-transfected with the TAT expressing plasmid pSZ90161:TAT. For both HPV18 expression plasmids, COS-7 cells are transfected using either DEAE-dextran or lipofection with Lipofectamine (BRL).

c) Expression in Insect Cells. The HPV18 L1-containing baculovirus transfer vector pAcUW51:HPV18 L1 is used to produce recombinant baculovirus (*Autographa californica*) by *in vivo*

- 33 -

homologous recombination. Epitope tagged HPV18 L1 is then expressed in Sf9 (*Spodoptera frugiperda*) insect cells grown in suspension culture following infection with the HPV18 - containing recombinant baculovirus.

5

EXAMPLE 19

Compounds that affect HPV18 activity may be detected by a variety of methods. A method of identifying compounds that affect HPV18 comprises:

10

- (a) mixing a test compound with a solution containing HPV18 to form a mixture;
- (b) measuring HPV18 activity in the mixture; and
- (c) comparing the HPV18 in the mixture to a standard.

15

Compounds that affect HPV18 activity may be formulated into pharmaceutical compositions. Such pharmaceutical compositions may be useful for treating diseases or conditions that are characterized by HPV18 infection.

20

EXAMPLE 20

DNA which is structurally related to DNA encoding HPV18 is detected with a probe. A suitable probe may be derived from DNA having all or a portion of the nucleotide sequence of Figure 1 or Figure 3, RNA encoded by DNA having all or a portion of the nucleotide sequence of Figure 1 or Figure 3 or degenerate oligonucleotides derived from a portion of the sequence of Figure 1 or Figure 3.

25

- 34 -

WHAT IS CLAIMED IS:

1. An isolated and purified DNA molecule which
encodes human papillomavirus type 18 or a functional derivative
5 thereof.
2. The isolated and purified DNA molecule of Claim 1
selected from the nucleotide sequence shown in Figure 1, the nucleotide
sequence shown in Figure 3, a functional derivative of the nucleotide
10 sequence shown in Figure 1, and a functional derivative of the nucleotide
sequence shown in Figure 3.
3. An expression vector for expression of the DNA
molecule of Claim 1 in a host.
15
4. An essentially purified protein encoded by the
DNA molecule of Claim 1.
5. An antibody immunologically reactive with the a
20 compound selected from the DNA molecule of Claim 1, RNA
complementary to the DNA molecule of Claim 1 or a protein encoded
by the DNA molecule of Claim 1.
6. A process for expression of a human
25 papillomavirus type 18 protein in a host, comprising:
 - (a) transferring the expression vector of Claim 4
into a suitable host; and
 - (b) culturing the host of step (a) under
30 conditions which allow expression of the
human papillomavirus type 18 protein from
the expression vector.

- 35 -

7. The process of Claim 6 wherein the protein is selected from HPV18 L1 protein, HPV18 L2 protein and combinations thereof.

5 8. A composition capable of inducing an immune response in a subject treated with the composition, the composition containing a compound selected from the DNA molecule of Claim 1, peptides encoded by the DNA molecule of Claim 1, RNA
10 complementary to the DNA molecule of Claim 1, or combinations thereof.

15 9. A vaccine for the prevention or treatment of human papillomavirus infection, the vaccine containing a compound selected from the group consisting of the DNA molecule of Claim 1, peptides encoded by the DNA molecule of Claim 1, RNA complementary to the DNA molecule of Claim 1, or combinations thereof.

20 10. A method for inducing immune responses against infection or disease caused by human papillomavirus which comprises introducing into an animal the DNA molecule of Claim 1, RNA complementary to the DNA molecule of Claim 1, protein encoded by the DNA molecule of Claim 1, or combinations thereof.

25 11. Virus-like particles comprised of recombinant L1 protein or recombinant L1 + L2 proteins of human papillomavirus 18, the virus-like particles being at least 60% pure.

30 12. The virus-like particles of Claim 11 wherein the recombinant L1 protein or the recombinant L1 + L2 proteins is produced in yeast.

13. A method of producing the virus-like particles of Claim 11, comprising:

- 36 -

- 5 (a) transforming yeast with a recombinant DNA molecule encoding papillomavirus L1 protein or papillomavirus L2 protein or papillomavirus L1 + L2 proteins;
- (b) cultivating the transformed yeast under conditions that permit expression of the recombinant DNA molecule to produce the recombinant papillomavirus protein; and
- 10 (c) isolating the recombinant papillomavirus protein to produce virus-like particles of Claim 1.
14. Recombinant papillomavirus protein produced by the method of Claim 13.
- 15 11. 15. A vaccine comprising the virus-like particles of Claim 11.
16. Pharmaceutical compositions comprising the virus-like particles of Claim 11.
- 20 17. A method of preventing papillomavirus infection comprising administering the vaccine of Claim 15 to a host.
- 25 18. A method for producing a yeast-derived recombinant papillomavirus capsid protein assembled into a virus-like particle, comprising:
- 30 a) cloning a papillomavirus gene that codes for at least one papillomavirus capsid protein into a vector;
- b) transferring the vector into a host cell to produce a recombinant host cell;
- c) cultivating the recombinant host cell under conditions that permit production of papillomavirus capsid protein; and

- 37 -

- d) purifying the papillomavirus capsid protein under conditions that permit formation of a virus-like particle.

5
Claim 18.

- 19. Virus-like particles produced by the method of

20. A method for inducing an immune response in an
animal comprising administering the virus-like particles of Claim 11 to an
10 animal.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
219

10 20 30 40 50 60
ATGGCTTTGTGGCGGCCTAGTGACAATACCGTATACCTTCCACCTCCTTCTGTGGCAAGA
1 M A L W R P S D N T V Y L P P P S V A R 20
70 80 90 100 110 120
GTTGTAAATACTGATGATTATGTGACTCGCACAAGCATATTTTATCATGCTGGCAGCTCT
21 V V N T D D Y V T R T S I F Y H A G S S 40
130 140 150 160 170 180
AGATTATTAACCTGTTGGTAATCCATATTTTAGGGTTCCTGCAGGTGGTGGCAATAAGCAG
41 R L L T V G N P Y F R V P A G G G N K Q 60
190 200 210 220 230 240
GATATTCCTAAGGTTTCTGCATACCAATATAGAGTATTTTCGGGTGCAGTTACCTGACCCA
61 D I P K V S A Y Q Y R V F R V Q L P D P 80
250 260 270 280 290 300
AATAAATTTGGTTTACCTGATAATAGTATTTATAATCCTGAAACACAACGTTTGTAGTGTGG
81 N K F G L P D N S I Y N P E T Q R L V W 100
310 320 330 340 350 360
GCCTGTGCTGGAGTGGAAATTGGCCGTGGTCAGCCTTTAGGTGTTGGCCTTAGTGGGCAT
101 A C A G V E I G R G Q P L G V G L S G H 120
370 380 390 400 410 420
CCATTTTATAATAAATTAGATGACACTGAAAGTTCCCATGCCGCTACGTCTAATGTTTCT
121 P F Y N K L D D T E S S H A A T S N V S 140
430 440 450 460 470 480
GAGGACGTTAGGGACAATGTGTCTGTAGATTATAAGCAGACACAGTTATGTATTTTGGGC
141 E D V R D N V S V D Y K Q T Q L C I L G 160
490 500 510 520 530 540
TGTGCCCCTGCTATTGGGGAACACTGGGCTAAAGGCACTGCTTGAAATCGCGTCCTTTA
161 C A P A I G E H W A K G T A C K S R P L 180
550 560 570 580 590 600
TCACAGGGCGATTGCCCCCTTTAGAACTTAAGAACACAGTTTGGGAAGATGGTGATATG
181 S Q G D C P P L E L K N T V L E D G D M 200
610 620 630 640 650 660
GTAGATACTGGATATGGTGCCATGGACTTTAGTACATTGCAAGATACTAAATGTGAGGTA
201 V D T G Y G A M D F S T L Q D T K C E V 220
670 680 690 700 710 720
CCATTGGATATTTGTCAGTCTATTTGTAAATATCCTGATTATTTACAAATGTCTGCAGAT
221 P L D I C Q S I C K Y P D Y L Q M S A D 240
730 740 750 760 770 780
CCTTATGGGGATTCCATGTTTTTTTGCTTACGACGTGAGCAGCTTTTGTAGGCATTTT
241 P Y G D S M F F C L R R E Q L F A R H F 260
790 800 810 820 830 840
TGGAATAGGGCAGGTACTATGGGTGACACTGTGCCTCAATCCTTATATATTAAAGGCACA
261 W N R A G T M G D T V P Q S L Y I K G T 280

FIG. 1A

2/7

850 860 870 880 890 900
GGTATGCGTGCTTCACCTGGCAGCTGTGTGTATTCTCCCTCTCCAAGTGGCTCTATTGTT
281 G M R A S P G S C V Y S P S P S G S I V 300
910 920 930 940 950 960
ACCTCTGACTCCCAGTTGTTTAATAAACCATATTGGTTACATAAGGCACAGGGTCATAAC
301 T S D S Q L F N K P Y W L H K A Q G H N 320
970 980 990 1000 1010 1020
AATGGTATCTGCTGGCATAATCAATTATTTGTTACTGTGGTAGATACCACTCGTAGTACC
321 N G I C W H N Q L F V T V V D T T R S T 340
1030 1040 1050 1060 1070 1080
AATTTAACAATATGTGCTTCTACACAGTCTCCTGTACCTGGGCAATATGATGCTACCAAA
341 N L T I C A S T Q S P V P G Q Y D A T K 360
1090 1100 1110 1120 1130 1140
TTTAAGCAGTATAGCAGACATGTTGAAGAATATGATTTGCAGTTTATTTTTTCAGTTATGT
361 F K Q Y S R H V E E Y D L Q F I F Q L C 380
1150 1160 1170 1180 1190 1200
ACTATTACTTTAACTGCAGATGTTATGTCCTATATTCATAGTATGAATAGCAGTATTTTA
381 T I T L T A D V M S Y I H S M N S S I L 400
1210 1220 1230 1240 1250 1260
GAGGATTGGAACCTTTGGTGTTCACCCCGCCAACTACTAGTTTGGTGGATACATATCGT
401 E D W N F G V P P P P T T S L V D T Y R 420
1270 1280 1290 1300 1310 1320
TTTGTACAATCTGTTGCTATTACCTGTCAAAAGGATGCTGCACCAGCTGAAAATAAGGAT
421 F V Q S V A I T C Q K D A A P A E N K D 440
1330 1340 1350 1360 1370 1380
CCCTATGATAAGTTAAAGTTTTGGAATGTGGATTTAAAGGAAAAGTTTTCTTTGGACTTA
441 P Y D K L K F W N V D L K E K F S L D L 460
1390 1400 1410 1420 1430 1440
GATCAATATCCCTTTGGACGTAAATTTTTGGTTCAGGCTGGATTGCGTCGCAAGCCCACC
461 D Q Y P L G R K F L V Q A G L R R K P T 480
1450 1460 1470 1480 1490 1500
ATAGGCCCTCGTAAACGTTCTGCTCCATCTGCCACTACGTCTTCTAAACCTGCCAAGCGT
481 I G P R K R S A P S A T T S S K P A K R 500
1510 1520
GTGCGTGACGTGCCAGGAAGTAA
501 V R V R A R K * 508

FIG. 1B

3/7

AMINO ACID VARIATIONS IN L1 PROTEIN OF HPV18					
	<u>AMINO ACID POSITION IN L1</u>				
	<u>30</u>	<u>88</u>	<u>283</u>	<u>323</u>	<u>338</u>
HPV18 PUBLISHED	P	T	P	V	P
HPV18 MERCK	R	N	R	I	R
#354 (CLINICAL INDIANA)	R	N	R	V	R
#556	R	N	R	V	R
#755	-	-	R	V	R
#697	-	-	R	V	R
#795	-	-	R	V	R
#23 (CLINICAL PENNSYLVANIA)	-	-	R	I	R

FIG.2

10 20 30 40 50 60
ATGGTATCCCACCGTGCCGCACGACGCAAACGGGCTTCGGTGACTGACTTATATAAAACA
1 M V S H R A A R R K R A S V T D L Y K T 20
70 80 90 100 110 120
TGTAACAATCTGGTACATGTCCATCTGATGTTGTTAATAAGGTAGAGGGCACCACGTTA
21 C K Q S G T C P S D V V N K V E G T T L 40
130 140 150 160 170 180
GCAGATAAAATAATGCAATGGTCAAGCCTTGATATTTTTGGGTGGACTTGGCATAGGT
41 A D K I L Q W S S L G I F L G G L G I G 60
190 200 210 220 230 240
ACTGGAAGTGGTACAGGGGGTCGTACAGGGTACATTCCATTGGGTGGGCGTTCCAATACA
61 T G S G T G G R T G Y I P L G G R S N T 80
250 260 270 280 290 300
GTTGTGGATGTCGGTCCTACACGTCCTCCAGTGGTTATTGAACCTGTGGGCCCCACAGAC
81 V V D V G P T R P P V V I E P V G P T D 100
310 320 330 340 350 360
CCATCTATTGTTACATTAATAGAGGACTCAAGTGTGTTACATCAGGTGCACCTAGGCCT
101 P S I V T L I E D S S V V T S G A P R P 120
370 380 390 400 410 420
ACTTTTACTGGCACGTCTGGGTTTGATATAACATCTGCTGGTACAACCTACACCTGCAGTT
121 T F T G T S G F D I T S A G T T T P A V 140
430 440 450 460 470 480
TTGGATATCACACCTTCGTCTACCTCTGTTTCTATTTCCACAACCAATTTTACCAATCCT
141 L D I T P S S T S V S I S T T N F T N P 160
490 500 510 520 530 540
GCATTTTCTGATCCGTCCATTATTGAAGTTCACAACTGGGGAGGTGTCAGGTAATGTA
161 A F S D P S I I E V P Q T G E V S G N V 180
550 560 570 580 590 600
TTTGTGGTACCCCTACATCTGGAACACATGGGTATGAAGAAATACCTTTACAAACATTT
181 F V G T P T S G T H G Y E E I P L Q T F 200
610 620 630 640 650 660
GCTTCTTCTGGTACGGGGGAGGAACCCATTAGTAGTACCCCATTCCTACTGTGCGGCGT
201 A S S G T G E E P I S S T P L P T V R R 220
670 680 690 700 710 720
GTAGCAGGTCCCCGCCTTTACAGTAGGGCCTACCAACAAGTGTCTGTGGCTAACCTGAG
221 V A G P R L Y S R A Y Q Q V S V A N P E 240
730 740 750 760 770 780
TTTCTTACACGTCCATCCTCTTTAATTACCTATGACAACCCGGCCTTTGAGCCTGTGGAC
241 F L T R P S S L I T Y D N P A F E P V D 260
790 800 810 820 830 840
ACTACATTAACATTTGAGCCTCGTAGTAATGTTCTGATTGAGATTTTATGGATATTATC
261 T T L T F E P R S N V P D S D F M D I I 280

FIG. 3A

SUBSTITUTE SHEET (RULE 26)

850 860 870 880 890 900
CGTTTACATAGGCCTGCTTTAACATCCAGGCGTGGTACTGTGCGCTTTAGTAGATTAGGT
281 R L H R P A L T S R R G T V R F S R L 300
910 920 930 940 950 960
CAAAGGGCAACTATGTTTACCCGTAGCGGTACACAAATAGGTGCTAGGGTTCACTTTTAT
301 Q R A T M F T R S G T Q I G A R V H F Y 320
970 980 990 1000 1010 1020
CATGATATAAGTCCTATTGCACCCTCCCCAGAATATATTGAACTGCAGCCTTTAGTATCT
321 H D I S P I A P S P E Y I E L Q P L V S 340
1030 1040 1050 1060 1070 1080
GCCACGGAGGACAATGGCTTGTTTGATATATATGCAGATGACATAGACCCTGCAATGCCT
341 A T E D N G L F D I Y A D D I D P A M P 360
1090 1100 1110 1020 1030 1040
GTACCATCGCGTCCTACTACCTCCTCTGCAGTTTCTACATATTCGCCCACTATATCATCT
361 V P S R P T T S S A V S T Y S P T I S S 380
1150 1160 1170 1180 1190 1200
GCCTCTTCCTATAGTAATGTAACGGTCCCTTTAACCTCCTCTTGGGATGTGCCTGTATAC
381 A S S Y S N V T V P L T S S W D V P V Y 400
1210 1220 1230 1240 1250 1260
ACGGGTCCTGATATTACATTACCACCTACTACCTCTGTATGGCCCATTTGTATCACCCACA
401 T G P D I T L P P T T S V W P I V S P T 420
1270 1280 1290 1300 1310 1320
GCCCCTGCCTCTACACAGTATATTGGTATACATGGTACACATTATTATTTGTGGCCATTA
421 A P A S T Q Y I G I H G T H Y Y L W P L 440
1330 1340 1350 1360 1370 1380
TATTATTTTATTCCTAAAAAGCGTAAACGTGTTCCCTATTTTTTTGCAGATGGCTTTGTG
441 Y Y F I P K K R K R V P Y F F A D G F V 460
GCGGCCTAG
461 A A * 463

FIG. 3B




FIG.5




FIG.4




FIG.6

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes Reference to PCT International Applications)

ATTORNEY'S DOCKET NUMBER

19424PC

NOV 2 1 1997
PATENT & TRADEMARK OFFICE
As below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

DNA ENCODING HUMAN PAPILLIOMAVIRUS TYPE 18

the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application

Serial No _____

on _____

and was amended

on _____ (if applicable).

☒ was filed as PCT international application

Number PCT/US96/03649

on _____

and was amended under PCT Article 19

on _____ (if applicable).

☐ is being filed as PCT international application, Attorney Docket No. _____

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (if PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued)
(Includes Reference to PCT International Applications)

ATTORNEY'S DOCKET NUMBER
19424PC

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS			STATUS (Check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED	
PCT APPLICATIONS DESIGNATING THE U.S.					
PCT APPLICATION NO.	PCT FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)	PATENTED	PENDING	ABANDONED

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number)

Send Correspondence to:

Joanne M. Giesser
Merck & Co., Inc.
P.O. Box 2000, 126 East Lincoln Avenue
Rahway, New Jersey 07065

Direct Telephone Calls to:
(name and telephone number)
732 594 3046

201	FULL NAME OF INVENTOR	FAMILY NAME HOFFMAN	FIRST GIVEN NAME KATHRYN	SECOND GIVEN NAME J.
	RESIDENCE & CITIZENSHIP	CITY COLLEGEVILLE	STATE OR FOREIGN COUNTRY PENNSYLVANIA 19426 PA	COUNTRY OF CITIZENSHIP UNITED STATES
	POST OFFICE ADDRESS	POST OFFICE ADDRESS P O BOX 2000	CITY RAHWAY	STATE & ZIP CODE/COUNTRY NEW JERSEY 07065 USA
202	FULL NAME OF INVENTOR	FAMILY NAME JANSEN	FIRST GIVEN NAME KATHRIN	SECOND GIVEN NAME U.
	RESIDENCE & CITIZENSHIP	CITY FORT WASHINGTON	STATE OR FOREIGN COUNTRY PENNSYLVANIA 19034 PA	COUNTRY OF CITIZENSHIP GERMANY
	POST OFFICE ADDRESS	POST OFFICE ADDRESS P O BOX 2000	CITY RAHWAY	STATE & ZIP CODE/COUNTRY NEW JERSEY 07065 USA
203	FULL NAME OF INVENTOR	FAMILY NAME NEEPER	FIRST GIVEN NAME MICHAEL	SECOND GIVEN NAME P.
	RESIDENCE & CITIZENSHIP	CITY COLLEGEVILLE	STATE OR FOREIGN COUNTRY PENNSYLVANIA 19426 PA	COUNTRY OF CITIZENSHIP UNITED STATES
	POST OFFICE ADDRESS	POST OFFICE ADDRESS P O BOX 2000	CITY RAHWAY	STATE & ZIP CODE/COUNTRY NEW JERSEY 07065 USA

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201 <i>Kathryn J. Hoffman</i>	SIGNATURE OF INVENTOR 202 <i>Kathrin Jansen</i>	SIGNATURE OF INVENTOR 203 <i>Michael Neeper</i>
DATE Nov. 4, 1997	DATE Nov. 6, 1997	DATE Nov. 3, 1997

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued) (Includes Reference to PCT International Applications)				ATTORNEY'S DOCKET NUMBER 19424PC
Continuation of Inventor Information				
204	FULL NAME OF INVENTOR	FAMILY NAME <u>JOYCE</u>	FIRST GIVEN NAME <u>JOSEPH</u>	SECOND GIVEN NAME <u>G.</u>
	RESIDENCE & CITIZENSHIP	CITY <u>LANSDALE</u>	STATE OR FOREIGN COUNTRY PENNSYLVANIA 19446 <u>PA</u>	COUNTRY OF CITIZENSHIP UNITED STATES
	POST OFFICE ADDRESS	POST OFFICE ADDRESS P O BOX 2000	CITY RAHWAY	STATE & ZIP CODE/COUNTRY NEW JERSEY 07065 USA
205	FULL NAME OF INVENTOR	FAMILY NAME <u>GEORGE</u>	FIRST GIVEN NAME <u>HUGH</u>	SECOND GIVEN NAME <u>A.</u>
	RESIDENCE & CITIZENSHIP	CITY <u>SCHWENKSVILLE</u>	STATE OR FOREIGN COUNTRY PENNSYLVANIA 19473 <u>PA</u>	COUNTRY OF CITIZENSHIP UNITED STATES
	POST OFFICE ADDRESS	POST OFFICE ADDRESS P O BOX 2000	CITY RAHWAY	STATE & ZIP CODE/COUNTRY NEW JERSEY 07065 USA
206	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
207	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
208	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
209	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.				
SIGNATURE OF INVENTOR 204 <i>Joseph L. Joyce</i>		SIGNATURE OF INVENTOR 205 <i>Hugh A. George</i>		SIGNATURE OF INVENTOR 206
DATE <i>November 3, 1997</i>		DATE <i>Nov 3, 1997</i>		DATE
SIGNATURE OF INVENTOR 207		SIGNATURE OF INVENTOR 208		SIGNATURE OF INVENTOR 209
DATE		DATE		DATE